

Geometry of spherically symmetric spacetime

Sebastiano Bernuzzi*

**Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, 07743, Jena, Germany*

(Dated: January 6, 2020)

Collection of general formulas for spherically symmetric spacetime in $n = 4$ dimensions. These calculations are preparatory to prove Birkhoff theorem.

GR class @ Jena FSU, WS 2019/20 <http://sbernuzzi.gitpages.tpi.uni-jena.de/gr/>

Follow notation of [1].

Assume the metric:

$$g = -e^{2\alpha(t,r)}dt^2 + e^{2\beta(t,r)}dr^2 + r^2d\Omega^2 \quad (1a)$$

with the 2-sphere metric

$$d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2 \quad (1b)$$

Christoffel symbols (only nonvanishing components):

$$\Gamma_{tt}^t = \partial_t\alpha \quad \Gamma_{tt}^r = e^{2(\alpha-\beta)}\partial_r\alpha \quad \Gamma_{\theta\theta}^r = -re^{-2\beta} \quad \Gamma_{\phi\phi}^\theta = -\sin\theta\cos\theta \quad (2a)$$

$$\Gamma_{tr}^t = \partial_r\alpha \quad \Gamma_{tr}^r = \partial_t\beta \quad \Gamma_{\phi\phi}^r = -re^{-2\beta}\sin^2\theta \quad \Gamma_{r\phi}^\phi = r^{-1} \quad (2b)$$

$$\Gamma_{rr}^t = e^{2(\beta-\alpha)}\partial_t\beta \quad \Gamma_{rr}^r = \partial_r\beta \quad \Gamma_{r\theta}^\theta = r^{-1} \quad \Gamma_{\theta\phi}^\phi = \frac{\cos\theta}{\sin\theta} \quad (2c)$$

Riemann tensor:

$$R_{rtr}^t = e^{2(\beta-\alpha)}(\partial_t^2\beta + (\partial_t\beta)^2 - \partial_t\alpha\partial_t\beta) + (\partial_r\alpha\partial_r\beta - \partial_r^2\alpha - (\partial_r\alpha)^2) \quad (3a)$$

$$R_{\theta t\theta}^t = -re^{-2\beta}\partial_r\alpha \quad R_{\phi t\phi}^t = -re^{-2\beta}\sin^2\theta\partial_r\alpha \quad R_{\theta r\theta}^t = -re^{-2\alpha}\partial_t\beta \quad (3b)$$

$$R_{\phi r\phi}^t = -re^{-2\alpha}\sin^2\theta\partial_t\beta \quad R_{\theta r\theta}^r = re^{-2\beta}\partial_r\beta \quad R_{\phi r\phi}^r = re^{-2\beta}\sin^2\theta\partial_r\beta \quad (3c)$$

$$R_{\phi\theta\phi}^\theta = (1 - e^{-2\beta})\sin^2\theta \quad (3d)$$

Ricci tensor:

$$R_{tt} = (\partial_t^2\beta + (\partial_t\beta)^2 - \partial_t\alpha\partial_t\beta) + e^{2(\alpha-\beta)}(\partial_r^2\alpha + (\partial_r\alpha)^2 - \partial_r\alpha\partial_r\beta + 2r^{-1}\partial_r\alpha) \quad (4a)$$

$$R_{rr} = -(\partial_r^2\alpha + (\partial_r\alpha)^2 - \partial_r\alpha\partial_r\beta - 2r^{-1}\partial_r\beta) + e^{2(\beta-\alpha)}(\partial_t^2\beta + (\partial_t\beta)^2 - \partial_t\alpha\partial_t\beta) \quad (4b)$$

$$R_{tr} = 2r^{-1}\partial_t\beta \quad R_{\theta\theta} = e^{-2\beta}(r\partial_r\beta - r\partial_r\alpha - 1) + 1 \quad R_{\phi\phi} = R_{\theta\theta}\sin^2\theta \quad (4c)$$